



IHC image of staining of rat amygdala

Neurotensin Antibody

Catalog #	20072	Product type	Primary antibodies
Lot #	1525001	Clonality	Polyclonal
Form	Lyophilized whole serum (100 µL)	Isotype	IgG
Host	Rabbit	Preservative	≤ 0.09% sodium azide
Reacts With	Cat, Dog, Frog, Human, Monkey, Mouse, Pigeon, Rat, Turtle	Antigen	Synthetic peptide (human) neurotensin coupled to bovine thyroglobulin (BTg) with glutaraldehyde.

INSTRUCTIONS

Preparation	<p>Do not reconstitute until ready to use since the product is most stable when lyophilized. The product does not need to be kept cooled during shipping; however, for long-term storage, store lyophilized antibody until ready to use at -15°C or lower. Reconstitute with 100 µL of distilled or deionized water. After reconstitution, use immediately or refrigerate at 2°–8°C. To avoid freeze/thaw cycles, dilute unused antibody with PBS or Tris buffer at a dilution no higher than 1/10, then aliquot and freeze at -15°C or lower.</p> <p>Refer to the Instruction Manual available online at www.immunostar.com for information on tissue preparation, immunostaining techniques, troubleshooting, and formulas.</p>
--------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

APPLICATION

IHC Quality Control	The ImmunoStar Neurotensin antiserum was quality control tested using standard immunohistochemical methods. The antiserum demonstrates significant labeling of rat amygdala using indirect immunofluorescent and biotin/avidin-HRP techniques. Staining is completely eliminated by pretreatment with 10 µg of Neurotensin per 1 mL of diluted antibody.
Tissue	Rat amygdala
Absorption Control	Neurotensin 10 ug/mL diluted serum.
Perfusion Fixation	<ul style="list-style-type: none"> Fixative: 4% paraformaldehyde in 0.1 M Phosphate buffer, pH 7.4; 500 mL over 20–30 min. Post Fixation: 1.5 hr at 4°C in 4% paraformaldehyde in 0.1 M phosphate buffer, pH 7.4.
Sections	10 µm cryostat or 50 µm vibratome
Tissue Incubation	18–24 hours at 2°–8°C
Detection System	Use Cy3 or Bn/Av-HRP according to manufacturers' directions.
Suggested Dilution	1/4,000–1/8,000 in PBS/0.3% Triton x-100 – Bn/Av-HRP immunohistochemistry

NOTES

Special Instructions	It is recommended that the researcher perform a primary antibody dilution series using our dilution recommendations as a guideline. Note that a change in the fixation or buffering system from our protocol may change the configuration of the protein which could alter the reactivity with the tissue tested.
Storage	After reconstitution, use immediately or refrigerate at 2°–8°C up to 2 days. For long-term storage, aliquot antibody and freeze at -15°C or lower. Avoid repeated freeze/thaw cycles.
Concentration	Not applicable. Antibody concentration is only relevant for purified antibodies.
Journal References	www.immunostar.com/publications

For Laboratory Reagent Use Only. Analytical and performance characteristics are not established.

ALL PRODUCTS ARE FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE RRID:AB_572254