



IHC image of neurons in rat brainstem.

## DBH (Dopamine-beta-Hydroxylase) Antibody

<b>Catalog #</b>	22806	<b>Product type</b>	Primary antibodies
<b>Lot #</b>	1905001	<b>Clonality</b>	Polyclonal
<b>Form</b>	Lyophilized whole serum (100 µL)	<b>Isotype</b>	IgG
<b>Host</b>	Rabbit	<b>Preservative</b>	≤ 0.09% sodium azide
<b>Reacts With</b>	Bird, Cat, Ferret, Finch, Guinea Pig, Hamster, Hatchefish, Human, Monkey, Mouse, Pig, Quail, Rat, Sea Snail, Sparrow, Starling, Steer, Turkey, Turtle, Zebra Finch	<b>Antigen</b>	DBH purified from bovine adrenal medulla.

### INSTRUCTIONS

<b>Preparation</b>	<p>Do not reconstitute until ready to use since the product is most stable when lyophilized. The product does not need to be kept cooled during shipping; however for long-term storage, store lyophilized antibody until ready to use at -15°C or lower. Reconstitute with 100 µL of distilled or deionized water. After reconstitution, use immediately or refrigerate at 2°–8°C or lower. To avoid freeze/thaw cycles, dilute unused antibody with PBS or Tris buffer at a dilution no higher than 1/10, then aliquot and freeze at -15°C or lower.</p> <p>Refer to the Instruction Manual available online at <a href="http://www.immunostar.com">www.immunostar.com</a> for information on tissue preparation, immunostaining techniques, troubleshooting, and formulas.</p>
--------------------	---

### APPLICATION

<b>IHC Quality Control</b>	The antibody has significant indirect immunofluorescence at 1/400 –1/800 and significant biotin-streptavidin/HRP staining at a 1/2,000–1/4,000 dilution in rat brain stem, cerebellum and adrenal medulla. Using western blot of purified DBH the antiserum detects a triplet at approximately 72–74 kD. Optimal dilution will vary depending upon fixation, labeling technique and/or detection system; therefore, a dilution series is recommended.
<b>Tissue</b>	Rat brainstem, cerebellum and adrenal medulla
<b>Perfusion Fixation</b>	<ul style="list-style-type: none"> <li>Fixative: 4% paraformaldehyde in 0.1M phosphate buffer, pH 7.4; 500 mL over 20 min.</li> <li>Post Fixation: 1.5 hour at 4°C in 4% paraformaldehyde in 0.1 M phosphate buffer, pH 7.4.</li> <li>Note: If needed, low levels of glutaraldehyde (0.1–0.3%) may be used in conjugation with paraformaldehyde.</li> </ul>
<b>Sections</b>	10 µm cryostat or 50 µm vibratome
<b>Tissue Incubation</b>	18–24 hours at 2°–8°C.
<b>Detection System</b>	Use Cy3 or Bn-SA/HRP reagents at dilutions recommended by the manufacturer.
<b>Suggested Dilution</b>	1/2,000–1/4,000 in PBS/0.3% Triton X-100 – Bn-SA/HRP immunohistochemistry

### NOTES

<b>Special Instructions</b>	It is recommended that the researcher perform a primary antibody dilution series using our dilution recommendations as a guideline. Note that a change in the fixation or buffering system from our protocol may change the configuration of the protein which could alter the reactivity with the tissue tested.
<b>Storage</b>	After reconstitution, use immediately or refrigerate at 2°–8°C up to 2 days. For long-term storage, aliquot and freeze at -15°C or lower. Avoid repeated freeze/thaw cycles.
<b>Concentration</b>	Not applicable. Antibody concentration is only relevant for purified antibodies.
<b>Journal References</b>	<a href="http://www.immunostar.com/publications">www.immunostar.com/publications</a>

For Laboratory Reagent Use Only. Analytical and performance characteristics are not established.

ALL PRODUCTS ARE FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE RRID:AB\_572229