

Anti-8-Hydroxy-2'-deoxyguanosine (8-OHdG) Monoclonal Antibody (N45.1)

- Catalog #:** 1) MOG-020P
2) MOG-100P
- Content:** 1) 20 μ g of IgG / vial. Lyophilized powder.
2) 100 μ g of IgG / vial. Lyophilized powder.
- Clone #:** N45.1
- Immunogen:** 8-Hydroxy-2'-deoxyguanosine (8-OHdG) conjugated keyhole limpet hemocyanin.
- Subclass:** Mouse IgG_{1(k)}, Prepared as ascite, and ammonium sulfate purified.
- Application:** Immunohistochemistry (Recommended concentration: 5~10 μ g/mL IgG).
ELISA
- Reconstitution:** 1) Reconstitute with 200 μ L of distilled water.
2) Reconstitute with 1000 μ L of distilled water.
- Buffer Concentration:** 100 μ g/mL IgG in 10mM Phosphate buffered saline, pH7.4 containing 1.0% BSA
- Specificity:** 19 analogues of 8-OHdG {guanosine (G), 7-methyl-G, 6-SH-G, 8-bromo-G, dA, dC, dT, dI, dU, dG, O6-methyl-dG, 8-OHdA, guanine (Gua), O6-methyl-Gua, 8-OHGua, uric acid, Urea, creatine, creatinine} demonstrate no cross-reactivity. Only 8-sulfhydryl-G and 8-OHG demonstrate minimal cross-reactivity (less than 1%). [ref.1]
- Storage:** Store at less than -20°C.
After reconstitution, store as frozen aliquots and avoid repeated freeze & thaw.
- Stability:** 5 years at -20°C.
- References:** Ref. 1) S.Toyokuni et al. Laboratory Investigation 76(3), p365-374 (1997)
Ref. 2) Y.Hattori et al. J.Invest.Dermatol. 107, p733-737 (1996)
Ref. 3) T.Tanaka et al. Laboratory Investigation 77(2), p145-155 (1997)

For research use only, not for diagnostic use.



JAPAN INSTITUTE FOR THE CONTROL OF AGING (JaICA)

710-1 Haruoka, Fukuroi, Shizuoka, 437-0122 Japan

TEL : +81-538-49-0125

FAX : +81-538-49-1267

WEB : <http://www.jaica.com>

E-mail : biotech@jaica.com