

REPRESENTATIVE DATASHEET

Goat anti-human Factor IX (FIX)

Peroxidase Conjugated Affinity-Purified IgG 0.1 mg

Product #: GAFIX-APHRP

Lot #: XXXX Expiry date: XXXX

Store at -10 to -20°C

1348 Sandhill Drive. Ancaster, Ontario, Canada L9G 4V5 905·304·9896 • 800·903·6020 • fax 905·304·9897

For Research Use Only. Not for use in diagnostic procedures.

Description of Factor IX

Factor IX (FIX, Christmas Factor) is a vitamin K-dependent glycoprotein produced in the liver. Plasma concentration of FIX is normally around 5 μ g/ml (87 nM) in plasma. The biological importance of F.IX is demonstrated in Haemophilia B (Christmas disease), an X-linked congenital bleeding disease resulting from a quantitative (low activity and low antigen) or qualitative (low activity and normal antigen) defect in FIX function.

In its proenzyme or zymogen form FIX is a single chain molecule of 55,000 daltons. It contains two EGF-like domains and an amino-terminal domain containing 12 γ -carboxy-glutamic acid (Gla) residues. These Gla residues allow FIX to bind divalent metal ions and participate in calcium-dependent binding interactions. The activation of FIX occurs by limited proteolysis in the presence of calcium by activated factor XI (FXIa) and/or by a complex of VIIa/tissue factor/phospholipid and activated Factor X between residues $\rm Arg^{146}\text{-}Ala^{147}$ and between $\rm Arg^{180}\text{-}Val^{181}$. The terminal activated product in either case is $\rm FIX^a{}_{\beta}$, a two-chain enzyme consisting of a heavy chain (28,000 daltons), a light chain (18,000 daltons) and an activation peptide product of 11,000 daltons. FIX can also be cleaved into inactive products by thrombin and by elastase.

The activity of $FIX^a{}_\beta$ in plasma is inhibited by antithrombin and this inhibition is accelerated 1000-fold in the presence of optimal concentrations of heparin $^{1-3}$.

REFERENCES and REVIEWS

- **1.** Lawson, JH, Mann KG; Cooperative Activation of Human F.IX by the Human Extrinsic Pathway of Coagulation; JBC 266 pp11317-11327, 10991.
- **2.** Enfield DL, Thompson AR; Cleavage and Activation of Factor IX by Serine Proteases; Blood 64, pp 821-831, 1984.
- **3.** Limentani SA, Furie BC, Furie B, in **Hemostasis and Thrombosis**, 3rd Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 94-108, J.B. Lippincott Co., Philadelphia PA, USA, 1994.

Product Specifications

Description:

Vial containing XXXX ml of affinity-purified IgG conjugated to horseradish peroxidase (HRP) through carbohydrate groups. Total protein is 0.1 mg.

Format:

APIgG-HRP conjugate as a clear, slightly red-brown liquid.

Host Animal:

Goat

Immunogen:

Human Factor IX purified from plasma.

Concentration:

APIgG-HRP concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient ($E^{1\%}_{280}$) of 14.

Buffer:

A buffered stabilizer solution containing 50% (v/v) glycerol.

Storage:

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use. Avoid exposure to Sodium azide as this is an inhibitor of peroxidase activity

Specificity:

Prior to conjugation, this antibody was specific for FIX as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of peroxidase labeled antibodies to FIX.

Rz Ratio (Reinheitszahl, A403/A280):

XXXX as determined spectrophotometrically.

Related Products:

Cat #: GAFIX-IG	Goat anti-human FIX, IgG from antiserum
Cat #: GAFIX-AP	Goat anti-human FIX, affinity-purified IgG
Cat #: GAFIX-HRP	Goat anti-human FIX, IgG-peroxidase
Cat #: SAFIX-IG	Sheep anti-human FIX, IgG from antiserum
Cat #: SAFIX-AP	Sheep anti-human FIX, affinity-purified IgG
Cat #: SAFIX-APHRP	Sheep anti-human FIX, APIgG-peroxidase
Cat #: FIX-EIA	Paired antibody set for ELISA of FIX, 5 x 96 wells
Cat #: FIX-DP	Human plasma deficient in FIX, immune depleted

Visit our site (www.affinitybiologicals.com) for details.

Limited Warranty: This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantibility or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by: **AFFINITY BIOLOGICALS INC.**

1348 Sandhill Drive Ancaster ON CANADA L9G 4V5

> Tel: (905) 304-9896 (800) 903-6020 Fax: (905) 304-9897

info@affinitybiologicals.com