

****REPRESENTATIVE DATASHEET****

Goat anti-human von Willebrand Factor

Whole IgG from antiserum

2.5 mg

Product #: GAVWF-IG

Lot #: XXXX

Expiry date: XXXX

Store at -10 to -20°C.

 1348 Sandhill Drive. Ancaster, Ontario, Canada L9G 4V5
 905-304-9896 • 800-903-6020 • fax 905-304-9897

For Research Use Only.

Not for use in diagnostic procedures.

Description of von Willebrand Factor

von Willebrand Factor (vWF, also previously referred to as Factor VIII related antigen) is a large adhesive protein produced in endothelial cells and megakaryocytes. There are two critical functions of vWF, the first being its involvement in the process of platelet adhesion and aggregation through interaction with platelet receptor glycoprotein Ib, the second being the binding and stabilization of Factor VIII (antihemophilic factor) for secretion and transport in plasma. The vWF precursor protein is synthesized with a 95,000 dalton propeptide (also known as vWF antigen-II), believed to be involved in the intracellular multimerization of the vWF subunits. The mature vWF multimers are then packed into storage organelles within the cell (Weibel-Palade bodies) after which the propeptide is cleaved and released. vWF circulates as multimers of disulphide linked 220,000 dalton subunits and the molecular weight of these multimers ranges from 0.5-20 million daltons.

The plasma concentration of vWF is typically 10 µg/ml, but increased levels are often observed in pregnancy and other conditions of physiological stress. von Willebrand's disease (vWD) is perhaps the most common inherited bleeding disorder in humans and is the result of either quantitative deficiencies of vWF (vWD Types I & III), or one of a number of qualitative disorders of vWF structure and function (vWD Type II).

REVIEW ARTICLES

1. Montgomery RR, Collier BS, in **Hemostasis and Thrombosis**, 3rd Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 134-168, J.B. Lippincott Co., Philadelphia PA, USA, 1994.

2. Sadler JE; von Willebrand Factor (Minireview); JBC 266:34, pp 22777-22780, 1991.

Product Specifications

Description:

Vial containing XXXX ml of whole IgG representing approximately 1 ml of antiserum. Total protein is 2.5 mg.

Format:

Whole IgG, clear liquid.

Host Animal:

Goat

Immunogen:

Human vWF purified from plasma.

Concentration:

 IgG concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient ($E_{280}^{1\%}$) of 13.4.

Buffer:

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

Storage:

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for vWF as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable for use as a source of antibodies to vWF.

Neutralizing activity:

Not determined

Species Cross Reactivity: (immunodiffusion vs. citrated plasma)

Human:	XXXX	Mouse:	XXXX	Rat:	XXXX
Rabbit:	XXXX	Pig:	XXXX	Dog:	XXXX

Related Products:

Cat #: **GAVWF-AP** Goat anti-human vWF, affinity purified IgG
 Cat #: **GAVWF-HRP** Goat anti-human vWF, IgG-peroxidase
 Cat #: **VWF-EIA** Paired antibody set for ELISA of vWF, 5 x 96 wells
 Cat #: **VWF-DP** Plasma deficient in vWF/FVIII, immune depleted
 Cat #: **VWF-LDP** Lyophilized plasma deficient in vWF/FVIII, immune depleted

 Visit our site (www.affinitybiologicals.com) for details.

Limited Warranty: This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantability or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by:
AFFINITY BIOLOGICALS INC.
 1348 Sandhill Drive
 Ancaster ON CANADA L9G 4V5
 Tel: (905) 304-9896
 (800) 903-6020
 Fax: (905) 304-9897
info@affinitybiologicals.com