

****REPRESENTATIVE DATASHEET****

Sheep anti-human Antithrombin (ATIII)

Affinity-Purified IgG

0.5 mg

Product #: SAAT-AP

Lot #: XXXX

Expiry date: XXXX

Store at -10 to -20°C.

1348 Sandhill Drive, Ancaster, Ontario, Canada L9G 4V5
905-304-9896 • 800-903-6020 • fax 905-304-9897

For Research Use Only.
Not for use in diagnostic procedures.

Description of Antithrombin (ATIII)

Antithrombin, also known as Antithrombin III (ATIII), is a member of the SERPIN family of proteinase inhibitors and the primary inhibitor of thrombin in plasma. It is produced in the liver and circulates in plasma at ~200 µg/ml (~3.5 µM). Antithrombin inhibits a broad spectrum of serine proteases including thrombin, activated forms of factor X, factor IX, factor XI, factor XII, as well as kallikrein, plasmin and urokinase. Enzyme inhibition by antithrombin occurs through proteolytic cleavage at Arg³⁸⁵-Ser³⁸⁶ and subsequent rapid formation of a stable, inactive 1:1 enzyme-antithrombin complex. Heparin has a profound accelerating effect on the inhibitory activity of antithrombin towards some enzymes. For example, the rate of inhibition of thrombin and activated factor X is increased 1000-fold in the presence of optimal concentrations of heparin, whereas heparin has relatively little effect on the inhibition rate of activated factor XI, activated factor XII and kallikrein. Antithrombin is a single chain molecule with a molecular weight of 59 kDa. Interaction with thrombin results in an SDS-stable thrombin-antithrombin complex of 96 kDa¹⁻³.

REFERENCES and REVIEWS

1. Damus PS, Rosenberg RD; Antithrombin - Heparin Cofactor; *Methods in Enzymology* 45, pp 653-669, 1976.
2. Harpel PC; Blood Proteolytic Enzyme Inhibitors: Their Role in Modulating Blood Coagulation and Fibrinolytic Enzyme Pathways; in *Hemostasis and Thrombosis*, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 738-747, J.B. Lippincott Co., Philadelphia PA, USA, 1982.
3. Griffith MJ, Noyes CM, Church FC; Reactive Site Peptide Structural Similarity between Heparin Cofactor II and Antithrombin III; *JBC*:260, pp 2218-2225, 1985.

Product Specifications

Description:

Vial containing XXXX ml of IgG purified by affinity-chromatography on immobilized ATIII. Total protein is 0.5 mg.

Format:

Affinity-purified IgG (APIgG), clear liquid.

Host Animal:

Sheep

Immunogen:

Human antithrombin purified from plasma.

Concentration:

APIgG concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient ($E^{1\%_{280}}$) of 13.4.

Buffer:

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

Storage:

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for antithrombin as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of enriched antibodies to antithrombin.

Neutralizing activity:

Not determined

Species Cross Reactivity: (immunodiffusion vs. citrated plasma)

Human:	XXXX	Mouse:	XXXX	Rat:	XXXX
Rabbit:	XXXX	Pig:	XXXX	Dog:	XXXX

Related Products:

Cat #: GAAT-IG	Goat anti-ATIII, whole IgG from antiserum
Cat #: GAAT-AP	Goat anti-ATIII, affinity-purified IgG
Cat #: SAAT-IG	Sheep anti-ATIII, whole IgG from antiserum
Cat #: SAAT-APHRP	Sheep anti-ATIII, APIgG-peroxidase
Cat #: ATIII-EIA	Paired antibody set for ELISA of ATIII, 5 x 96 wells
Cat #: TAT-EIA	Paired antibody ELISA of Thrombin-ATIII complexes
Cat #: AT-NU10	Paired antibody set for ATIII neutralization in plasma
Cat #: ATIII-DP	Human plasma deficient in ATIII, immune depleted
Cat #: ATHC-DP	Human ATIII & HCII def plasma, immune depleted

Visit our site (www.affinitybiologicals.com) for details.

Limited Warranty: This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantability or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by:
AFFINITY BIOLOGICALS INC.
1348 Sandhill Drive
Ancaster ON CANADA L9G 4V5
Tel: (905) 304-9896
(800) 903-6020
Fax: (905) 304-9897
info@affinitybiologicals.com