


# Datasheet




Mouse mAb to **CD19**  
Clone **FR10A7**  
Isotype **IgM-κ**

## Source

A BALB/c mouse was immunized with RAJI cells.  
Fusion partner: X63AG8.653.

## Specifications

FR10A7 is specific for the antigen CD19. This antigen has a MW of 120 kDa and contains a 280 residue extracellular domain and a 240 residue cytoplasmic domain. It is a critical signal transduction molecule that regulates B-lymphocyte development, activation, and differentiation. It plays a dominant role in establishing signalling thresholds for antigen receptors and other surface receptors on B-lymphocytes. This antigen is lost upon terminal differentiation to plasma cells.



**Figure 1:** Human PBL stained for CD19 (FACS).

## Species reactivity

Positive: human.

## Applications

FR10A7 can be used for immunophenotyping of leukemia and malignant cells in frozen tissue, B-lymphocyte detection in peripheral blood, B-cell localization in tissues and B-lymphocyte purification by immunoabsorption methods.

Flow cytometry	Frozen sections	Immunofluorescence
+	+	+

## Format

Produced in tissue culture, contains no host Ig. Antibodies are affinity purified and presented in PBS with 0,02% sodium azide.

Stored at 4°C-8°C, shelf life is at least 24 months after purchase.

## Dilution advice

- Flow cytometry (0,5-1,0 µg/million cells in 0,1 ml).
- Immunofluorescence (0,5-1,0 µg/ml).
- Immunohistology (1-2 µg/ml for 30-60 min at RT; information on a suitable antigen retrieval method for staining of formalin-fixed tissues is unavailable to date).

## Positive control

Primate PBLs.

## References

- Deaglio, S., et. al. *J Immunol.* **160(1)**: 395-402 (1998).