Anti PACAP27 (13-27) (Human) Serum Cat. No. Y051 Lot No. 082271206

Description: This antiserum was raised in a rabbit by immunization with a carrier free synthetic PACAP27 (13-27) (human) peptide. The product vial contains 50 μ L of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with the C-terminal portion of PACAP27 (human).

Immunogen: Synthetic PACAP27 (13-27) (human) peptide, carrier free

13

Host: Rabbit

Amino Acid Sequence of PACAP27 (13-27) (human)¹⁾:

HSDGIFTDSY SRYRKOMAVK KYLAAVL-NH2

Product Form: Lyophilized unpurified serum

Size: 50 µL

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).

Storage: The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezingthawing should be avoided.

Suggested Working Dilution Range: 1:2,000-20,000 (final dilution ~1:140,000) for radioimmunoassay; 1: 1,000-6,000 for immunohistochemistry (frozen or paraffin sections). Optimal dilution should be determined by each laboratory for each application.

Specificity (based on radioimmunoassay): PACAP27 (13-27) (human) 100%, PACAP27 (human) 149.9%, PACAP27 (8-27) (human) 112.9%, PACAP27 (16-27) (human) 54.9%, PACAP (22-27) (human) 3.3%, PACAP38 (human)<0.1%, PACAP38 (1-15) (human) 0%, VIP 0%, glucagon 0%, PHI (human) 0%, secretin (human) 0%

Positive Control (immunohistochemistry): Rat colon

Species Tested: Rat

REFERENCES:

1) A. Miyata, A. Arimura et al., Isolation of a neuropeptide corresponding to the N-terminal 27 residues of the pituitary adenylate cyclase activating polypeptide with 38 residues (PACAP38), Biochemistry Biophysics Research Communication 170:643-648, 1990

FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

Yanairora Institute Inc.

2480-1 Awakura, Fujinomiya, Shizuoka, 418-0011 Japan FAX:+81-544-22-2770

TEL: +81-544-22-2771

For more infomation, please visit our internet website: http://www.yanaihara.co.jp or contact us to:/ask@yanaihara.co.jp