

Anti Met-Enkephalin-Arg⁶-Gly⁷-Leu⁸ Serum

Cat. No. Y140 Lot No. 017171004

Description: This antiserum was raised in a rabbit by immunization with a *Ascaris suilla* protein conjugate of synthetic met-enkephalin-arg⁶-gly⁷-leu⁸ peptide. The product vial contains 50 μ L of the titled antiserum obtained by lyophilizing its 0.001M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with met-enkephalin-arg⁶-gly⁷-leu⁸.

Immunogen: Synthetic met-enkephalin-arg⁶-gly⁷-leu⁸-*Ascaris suilla* protein conjugate **Host:** Rabbit

Amino Acid Sequence of Met-Enkephalin-Arg⁶-Gly⁷-Leu⁸ ¹⁾: YGGFM RGL

Product Form: Lyophilized unpurified serum **Size:** 50 μ L

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).

Storage: The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

Suggested Working Dilution Range: 1:8,000 (final dilution~1:56,000) for radioimmunoassay; 1: 1,000-5,000 for immunohistochemistry (frozen or paraffin sections). Optimal dilution should be determined by each laboratory for each application.

Specificity (based on radioimmunoassay): Met-enkephalin-arg⁶-gly⁷-leu⁸ 100%, phe⁴-met⁵-arg⁶-gly⁷-leu⁸ < 0.001%, met-enkephalin 0%, leu-enkephalin 0%, α -endorphin 0%, β -endorphin 0%,

Positive Control (immunohistochemistry): Rat adrenal gland and colon

Species Tested: Bovine, guinea pig, dog, cat, hamster, human, porcine, monkey, rat^{2,3,4,5)}

REFERENCES:

- 1) P.L. Kilpatrick, B.N. Jones et al., Identification of the octapeptide [Met] enkephalin-Arg⁶-Gly⁷-Leu⁸ in extracts of bovine adrenal medulla. Biochemical Biophysical Research Communications 103:697-705, 1981
- 2) S. Kobayashi, N. Yanaihara et al., Immunocytochemical demonstration of the co-storage of noradrenaline with met-enkephalin-arg⁶-phe⁷ and met-enkephalin-arg⁶-gly⁷-leu⁸ in the carotid body chief cells of the dog. Archivum Histologicum Japonicum 46: 713-722, 1983
- 3) S. Kobayashi, N. Yanaihara et al., Met-enkephalin-arg-gly-leu-like immunoreactivity in adrenal chromaffin cells and carotid body chief cells of the dog and monkey. Biomedical Research 4: 201-210, 1983
- 4) K. Nihei, T. Fujita et al., Preproenkephalin A occurs in the entero-chromaffin (EC) cells of the porcine intestine: an immunocytochemical study using antisera to met-enkephalin-arg⁶-gly⁷-leu⁸ and to serotonin. Biomedical Research 4: 393-398, 1983
- 5) S. Kobayashi, N. Yanaihara et al., An immunohistochemical study on the co-storage of met-enkephalin-arg⁶-gly⁷-leu⁸ and met-enkephalin-arg⁶-phe⁷ with adrenaline and/or noradrenaline in the adrenal chromaffin cells of the rat, dog, and cat. Biomedical Research 4: 433-442, 1983

FOR RESEARCH LABORATORY USE ONLY
DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM



Yanaihara Institute Inc.

2480-1 Awakura, Fujinomiya, Shizuoka, 418-0011 Japan

TEL: +81-544-22-2771

FAX: +81-544-22-2770

For more information, please visit our internet website: <http://www.yanaihara.co.jp> or contact us to: ask@yanaihara.co.jp