

**(Suc-LLVY)2R110**

Catalog number: 13451

Unit size: 1 mg

Component	Storage	Amount
(Suc-LLVY)2R110	Freeze (<-15 °C), Minimize light exposure	1 mg

**OVERVIEW**

(Suc-LLVY)2R110 is a sensitive fluorogenic substrate for 20S proteasome, calpains and other chymotrypsin-like proteases. The non-fluorescent substrate generates a bright green fluorescent rhodamine 110 product that has an emission spectra that can be easily detected with a FITC filter set. This rhodamine 110 substrate is much more sensitive than the AMC-, AFC- or 4-nitroaniline-based substrates. The most common form of the proteasome is known as the 26S proteasome that contains one 20S core particle structure and two 19S regulatory caps. All 20S particles consist of four stacked heptameric ring structures that are themselves composed of two different types of subunits; alpha subunits are structural in nature, whereas beta subunits are predominantly catalytic. The outer two rings in the stack consist of seven alpha subunits each, which serve as docking domains for the regulatory particles and the alpha subunits N-termini form a gate that blocks unregulated access of substrates to the interior cavity. The inner two rings each consist of seven beta subunits and contain the protease active sites that perform the proteolysis reactions.

**AT A GLANCE**

**Important** Following protocol only provides a guideline, and should be modified according to your specific needs.

Product Number	Indicators	Excitation	Emission
13451	(Suc-LLVY)2R110	498 nm	520 nm
13453	Suc-LLVY-AMC	351 nm	430 nm
13455	(Ac-ANW)2R110	498 nm	520 nm
13465	(Ac-KQL)2R110	498 nm	520 nm
13466	(Z-LLE)2R110	498 nm	520 nm
13467	(Ac-PAL)2R110	498 nm	520 nm
13468	(Ac-WLA)2R110	498 nm	520 nm

**KEY PARAMETERS**

Instrument:	Fluorescence microplate reader
Excitation:	498 nm
Emission:	520 nm
Recommended plate:	Solid black

**PREPARATION OF STOCK SOLUTIONS**

Unless otherwise noted, all unused stock solutions should be divided into single-use aliquots and stored at -20 °C after preparation. Avoid repeated freeze-thaw cycles.


1. (Suc-LLVY)2R110 stock solution:  
Prepare a 5 to 10 mM stock solution in DMSO.

**PREPARATION OF WORKING SOLUTION**

<u>Component</u>	<u>Volume</u>
Substrate stock solution (10 mM)	25 µL
DTT (1M)	100 µL
EDTA (100 mM)	400 µL
Hepes Buffer (25 mM), pH = 7.4	10 mL
Total volume	10.53 mL

**SAMPLE EXPERIMENTAL PROTOCOL**

1. Mix equal volume of the proteasome standards or samples with 2X fluorescent proteasome substrate assay solution, and incubate the solutions at room temperature for at least 1 hour.
2. Monitor the fluorescence using fluorescent microplate readers at 498/520 nm.

**EXAMPLE DATA ANALYSIS AND FIGURES**


**Figure 1.** Chemical structure for (Suc-LLVY)2R110

**DISCLAIMER**

AAT Bioquest provides high-quality reagents and materials for research use only. For proper handling of potentially hazardous chemicals, please consult the Safety Data Sheet (SDS) provided for the product. Chemical analysis and/or reverse engineering of any kit or its components is strictly prohibited without written permission from AAT Bioquest. Please call 408-733-1055 or email info@aatbio.com if you have any questions.