

## iFluor™ 790 goat anti-mouse IgG (H+L) \*Cross Adsorbed\*

Catalog number: 16587, 16790 Unit size: 200 ug, 1 mg

**Product Details** 

Storage Conditions 2-6°C and kept from light. To extend the shelf-life of this product, add an equal volume of

glycerol to make a final concentration of approximately 50% glycerol and store at -20°C.

Expiration Date 12 months upon receiving

Concentration 1 mg/mL

Formulation PBS, 2 mg/mL BSA

**Unit Details** 

Unit 16587 (200 ug) 16790 (1 mg)

Reconstitution Volume 200 uL ddH<sub>2</sub>O 1 mL ddH<sub>2</sub>O

**Antibody Properties** 

Species Reactivity Mouse

Class Secondary

Clonality Polyclonal

Host Goat

**Chemical Properties** 

Molecular Weight ~150000

**Biological Properties** 

Stabilizer None

Appearance Green solid

Preparation Goat anti-mouse IgG (H+L) is produced in goat with pooled total mouse IgG, and affinity purified

with mouse IgG coupled beads. The purified IgG has a minimal cross-reaction to human, horse, rabbit and bovine IgG. The antibody is conjugated with iFluor™ 790 under optimal condition.

Immunofluorescence (IF), Flow Cytometry (FACS)

Soluble In Water

**Spectral Properties** 

Application

Conjugate iFluor™ 790

Excitation Wavelength 787 nm

Emission Wavelength 812 nm

## **Applications**

AAT Bioquest's iFluor™ dyes are optimized for labeling proteins, in particular, antibodies. These dyes are bright, photostable and have minimal quenching on proteins. They can be well excited by the major laser lines of fluorescence instruments (e.g., 350, 405, 488, 555 and 633 nm). iFluor™ 790 goat anti-mouse IgG (H+L) conjugate has IR fluorescence excitation and emission maxima of ~780 nm and ~810 nm respectively. These spectral characteristics make them an excellent alternative to IRDye® 800 goat anti-mouse IgG (H+L) conjugate (IRDye® is the trademark of Li-COR).