

**iFluor™ 810 goat anti-mouse IgG (H+L)
*Cross Adsorbed***Catalog number: 48006, 48007
Unit size: 200 µg, 1 mg**Product Details**

Storage Conditions	2-6°C and kept from light. To extend the shelf-life of this product, add an equal volume of glycerol to make a final concentration of approximately 50% glycerol and store at -20°C.
Expiration Date	12 months upon receiving
Concentration	1 mg/mL
Formulation	PBS, 2 mg/mL BSA

Unit Details

Unit	48006 (200 µg)	48007 (1 mg)
Reconstitution Volume	200 µL ddH ₂ O	1 mL ddH ₂ O

Antibody Properties

Species Reactivity	Mouse
Class	Secondary
Clonality	Polyclonal
Host	Goat

Chemical Properties

Molecular Weight	~150000
------------------	---------

Biological Properties

Stabilizer	None
Appearance	Green solid
Preparation	Goat anti-mouse IgG (H+L) is produced in goat with pooled total mouse IgG, and affinity purified with mouse IgG coupled beads. The purified IgG has a minimal cross-reaction to human, horse, rabbit, human and bovine IgG. The antibody is conjugated with iFluor™ 810 under optimal condition.
Application	Flow Cytometry (FACS), ELISA, HC, Western Blot
Soluble In	Water

Spectral Properties

Conjugate	iFluor™ 810
-----------	-------------

Excitation Wavelength	811 nm
Emission Wavelength	822 nm

Applications

AAT Bioquest's iFluor™ dyes are optimized for labeling proteins, in particular, antibodies. These dyes are bright, photostable, and have minimal quenching on proteins. They can be well excited by the major laser lines of fluorescence instruments (e.g., 350, 405, 488, 532-561, 633-647, and 808 nm). iFluor™ 810 goat anti-mouse IgG (H+L) conjugate has fluorescence excitation and emission maxima of 811 nm and 822 nm, respectively. These unique spectral characteristics makes iFluor™ 810 goat anti-mouse IgG (H+L) conjugates ideal for various NIR imaging applications, including Western blotting, ELISA, protein arrays, tissue section imaging, and *in vivo* imaging.